

GENERALISED THURSTON-BENNEQUIN INVARIANTS FOR REAL ALGEBRAIC SURFACE SINGULARITIES

FERİT ÖZTÜRK

ABSTRACT. A generalised Thurston-Bennequin invariant for a $\mathbb{Q}$ -singularity of a real algebraic variety is defined as a *linking form* on the homologies of the real link of the singularity. The main goal of this paper is to present a method to calculate the linking form in terms of the very good resolution graph of a real normal unibranch singularity of a real algebraic surface. For such singularities, the value of the linking form is the Thurston-Bennequin number of the real link of the singularity. As a special case of unibranch surface singularities, the behaviour of the linking form is investigated on the Brieskorn double points $x^m + y^n \pm z^2 = 0$.

1. INTRODUCTION

The link of a real algebraic surface singularity is a contact 3-manifold with the canonical contact structure the set of complex tangencies. Furthermore, the real link in this contact manifold is a Legendrian link. The *linking form* is defined as a rational valued form on the first homology of the real link whenever the rational linking number is well-defined in the link of the singularity, i.e. when the link is a $\mathbb{Q}$ -homology sphere (see Section 2.2 or [9] for the definition of the linking form). For such a link, the linking form, when evaluated on a pair of classes of knots of the real link, gives the rational linking number of the knots with contact framing. In the case when the real link is connected, i.e. the singularity is unibranch, the outcome of the linking form is by definition the Thurston-Bennequin number.

Therefore, one can consider the linking form as a *generalised* Thurston-Bennequin invariant. It is generalised in the sense that (i) it is defined for a *link* in a contact manifold, (ii) the knots of the link are not necessarily integrally null-homologous but they bound rationally, (iii) it is rational valued.

A method to compute this linking form has already been presented for surface singularities $f(x, y) \pm z^2 = 0$ ($f(x, y)$ reduced) using real deformations of the singularity [10].

The main result of this paper is Theorem 15, Section 5, which presents a method to calculate the generalised Thurston-Bennequin number of the connected real link of a normal real algebraic surface $\mathbb{Q}$ -singularity using a very good resolution graph of the singularity with the condition that the singularity is numerically Gorenstein. In particular, Theorem 15 is valid for all the unibranch hypersurface singularities in $\mathbb{C}^3$. Furthermore, we give an example which exhibits how the method works algorithmically for singularities with more than one real branch.

In the proof of Theorem 15, we make use of the fact that the generalised Thurston-Bennequin number of the connected real link of a surface singularity gives by definition the self-intersection of the oriented real part of the surface in the complex surface relative to the link of the singularity. This is not valid anymore if the real part is non-oriented. Theorem 15 determines how much the generalised Thurston-Bennequin number differs from the self-intersection of the real part. This difference is a non-integer rational number in general.

In the second part of the article, as a special case of hypersurface singularities, for which Theorem 15 is valid, we deduce some properties for the Thurston-Bennequin numbers of the Brieskorn double points $x^m + y^n \pm z^2 = 0$ with m, n relatively prime. We first express the conditions under which the

Thurston-Bennequin numbers become integer (see Corollary 18 and Corollary 20, Section 6). Then we investigate the behaviour of the Thurston-Bennequin numbers with changing m and n (see Theorems 21, 22 and 24, Section 6). These theorems show that one can calculate the Thurston-Bennequin numbers for large m and n in terms of the Thurston-Bennequin numbers corresponding to sufficiently small m and n .

The main approach in the proofs is to use Theorem 15. In fact, Corollary 18 is a direct consequence of Theorem 15. We prove Corollary 20 and Theorem 21 by observing the change in the very good resolution graph of the singularity when m is held fixed and n is varied (see Section 3) and using the results of Section 4.

The proofs of Theorems 22 and 24, Section 6, do not involve resolution graphs. Rather, we construct new real algebraic surfaces as branched double coverings of appropriate weighted homogeneous surfaces; these new algebraic surfaces contain at the same time both singularities involved in each theorem.

Notation: In this article, the manifold $\overline{X}$ is usually the quotient of the manifold X by complex conjugation except in the case $\overline{\mathbb{CP}^2}$ which is $\mathbb{CP}^2$ with reverse orientation.

Acknowledgments. I would like to thank Sergey Finashin (METU, Ankara) for having introduced me the problem, for his invaluable guidance and his inspiration and to Yıldıray Ozan (METU, Ankara) for many stimulating discussions. In addition, I would like to thank CMAT, Ecole Polytechnique, Palaiseau where I completed this work as a postdoc.

2. PRELIMINARY NOTIONS AND DEFINITIONS

2.1. The real branches of a real surface $\mathbb{Q}$ -singularity and the linking form. Let $\mathbb{C}X$ be the complex point set of a real algebraic surface with complex conjugation $\text{conj} : \mathbb{C}X \rightarrow \mathbb{C}X$; let $\mathbb{R}X$ denote the set of the real points of $\mathbb{C}X$ and $\overline{X}$ denote the quotient $\mathbb{C}X/\text{conj}$. Let x be an isolated singularity of $\mathbb{C}X$ and $\mathbb{C}U$ be a small compact cone-like neighbourhood of x [16].

It is known that the boundary $\mathbb{C}M = \partial\mathbb{C}U$ is a 3-manifold with a canonical contact structure determined by the distribution of complex lines that exist uniquely in the tangent space over each point of $\mathbb{C}M$ [22]. Furthermore, since the real link $\mathbb{R}M$ is tangent to the unique complex line at each point, $\mathbb{R}M$ is a Legendrian link in $\mathbb{C}M$. Although $\mathbb{C}M$ is not a complex manifold, we use the notation with prefix $\mathbb{C}$ to emphasize that $\mathbb{C}M$ is fixed under conj of $\mathbb{C}X$. With the same idea in mind, $\mathbb{R}M$ denotes the subset of $\mathbb{C}M$ fixed *pointwise* by conj . Finally $\overline{M}$ will denote $\partial\overline{U}$.

A singular point $x \in \mathbb{C}X$ is called a *topologically rational singularity* (or a *$\mathbb{Q}$ -singularity*) if its link $\mathbb{C}M$ is a $\mathbb{Q}$ -homology sphere. Point x is called a *$\overline{\mathbb{Q}}$ -singularity* if $\overline{M}$ is a $\mathbb{Q}$ -homology sphere. Let $x \in \mathbb{C}X$ be a $\mathbb{Q}$ -singularity and let σ denote the number of connected components of the real link $\mathbb{R}M$. Each connected component of $\mathbb{R}M$ is called a *real branch* of $\mathbb{C}X$ at x . If $\sigma = 1$, i.e. if $\mathbb{R}M$ is a knot, then we will say that x is a *real unibranch singularity*. For example, the Brieskorn singularity $\{x^m + y^n + z^d = 0\}$, $\gcd(m, n, d) = 1$ is an isolated $\mathbb{Q}$ -singularity which is a real unibranch singularity.

Let $\gamma_1, \dots, \gamma_\sigma$ be the connected components of $\mathbb{R}M$ and ξ_i be a *Thurston-Bennequin framing* of γ_i in $\mathbb{C}M$, i.e. a choice of trivialisation of a tubular neighbourhood of γ_i in $\mathbb{C}M$. Let us denote by $\gamma_i^{\xi_i}$ the knot obtained by a small shift of γ_i in the direction of ξ_i . Consider the bilinear form

$$\text{tb}_x : H_1(\mathbb{R}M; \mathbb{Q}) \times H_1(\mathbb{R}M; \mathbb{Q}) \rightarrow \mathbb{Q}$$

defined on the generators by:

$$\text{tb}_x([\gamma_i], [\gamma_j]) = \text{lk}(\gamma_i, \gamma_j^{\xi_j}) \text{ in } \mathbb{C}M$$

and extended linearly over $H_1(\mathbb{R}M; \mathbb{Q})$ (see [9] or [10]). Here lk is the rational linking number in $\mathbb{C}M$. Note that if $i = j$, then $\text{tb}_x([\gamma_i], [\gamma_i]) = \text{tb}(\gamma_i)$, the rational Thurston-Bennequin number of the Legendrian knot [3]. The form tb_x is called the *linking form*.

It is essential to note the following: Let σ_i denote the component of closure of $\mathbb{R}U - \{x\}$ containing γ_i ; the real surface σ_i is a cone over γ_i topologically. Consider a tangent vector field ν_i on σ_i directed outward along γ_i . Then $\sqrt{-1} \cdot \nu_i$ coincides with ξ_i , the Thurston-Bennequin framing, proving that in case σ_i and σ_j are oriented, the form tb_x evaluated on $[\gamma_i]$ and $[\gamma_j]$ is nothing but the rational intersection of σ_i and σ_j in $\mathbb{C}U$ relative to the boundary, i.e. :

$$(1) \quad \text{tb}_x([\gamma_i], [\gamma_j]) = \langle \sigma_i, \sigma_j \rangle_{(\mathbb{C}U, \mathbb{C}M)}.$$

Similarly as above, for a $\overline{Q}$ -singularity, one can define a bilinear form $\overline{\text{tb}}_x$ on $H_1(\mathbb{R}M; \mathbb{Q})$ by letting

$$\overline{\text{tb}}_x([\gamma_i], [\gamma_j]) = \text{lk}(q(\gamma_i), q(\gamma_j^{\xi_j})) \text{ in } \overline{M} = q(\mathbb{C}M)$$

where $q : \mathbb{C}X \rightarrow \overline{X}$ is the quotient map. For a $\overline{Q}$ -singularity, tb_x and $\overline{\text{tb}}_x$ satisfy [10]:

$$(2) \quad \overline{\text{tb}}_x = 2 \text{tb}_x.$$

The linking form tb_x corresponding to a singularity with σ real branches is represented by a $\sigma \times \sigma$ matrix. We are going to denote by $[\text{tb}_x]$ the $\sigma \times \sigma$ matrix in the basis $\gamma_1, \dots, \gamma_\sigma$. We will not mention γ_i s whenever there is no confusion and we will also allow permuted indexing. The matrix $[\text{tb}_x]$ of a real unibranch singularity x is a 1×1 matrix and in that case, tb_x will be called *the Thurston-Bennequin number* of the real unibranch singularity x as well.

An algorithmic method was introduced in [10], Theorem B, to compute the Thurston-Bennequin numbers of the real unibranch singularities $f(x, y) \pm z^2 = 0$ where $f(x, y)$ is a real polynomial with an isolated real singularity. This method makes use of the diagrams of Gusein-Zade [13] and A'Campo [1] which are obtained by small real generic deformations of the singularity $f(x, y) = 0$ in $\mathbb{C}^2$.

A more general approach for computation of the Thurston-Bennequin number associated to a normal unibranch Q -singularity was partly announced in [10] using a resolution of the singularity. Below, Theorem 15, Section 5, offers a method to compute the Thurston-Bennequin number of a real normal unibranch Q -singularity using a very good resolution graph of the singularity. In the case the real part of the ambient real algebraic surface is oriented, Theorem 15 gives Equation 1.

2.2. Resolution graphs. Let $\mathbb{C}X$ be a normal complex surface and $\rho : \widetilde{\mathbb{C}X} \rightarrow \mathbb{C}X$ be a resolution of $\mathbb{C}X$. The preimage $E = \rho^{-1}(\text{sing}(\mathbb{C}X)) \subset \widetilde{\mathbb{C}X}$ is called the exceptional set and each irreducible curve in the exceptional set is called an exceptional curve of the resolution ρ . Let the number of exceptional curves in E be s ; then the exceptional curves of E will be denoted by $E_1, \dots, E_s$.

The resolution ρ is said to be *good* if the exceptional curves are smoothly embedded and any two intersecting exceptional curves intersect at normal crossings. A *very good* resolution is a good resolution with any pair of exceptional curves having at most one common point. A good and a very good resolution always exist for any analytic surface (cf. [15] or [7]).

The dual graph of a resolution of a surface singularity is constructed by representing each exceptional curve by a vertex and connecting two vertices by an edge if the corresponding exceptional curves intersect. It is well known that the dual graph Γ of a good resolution of a normal surface singularity is connected. The s vertices $e_1, \dots, e_s$ of Γ correspond to $E_1, \dots, E_s$ respectively. Each vertex e_i ($1 \leq i \leq s$) is endowed with a pair of integers: the self intersection n_i and the genus g_i of E_i . Two vertices e_i and e_j are connected by n_{ij} edges where n_{ij} is the number of points in $E_i \cap E_j$.

From now on, let us assume that Γ is the dual graph of a very good resolution $\rho : \widetilde{\mathbb{C}X} \rightarrow \mathbb{C}X$ and also that Γ is a tree.

If a vertex e of Γ is connected to more than two vertices, then e is called a *rupture vertex*. A vertex adjacent to one and only one vertex of Γ is called a *terminal vertex*. The *star* of a subgraph σ of Γ is a subgraph of Γ consisting of σ and all the edges connected to any vertex of σ . The star of σ is denoted by $\text{star}(\sigma)$ (see Figure 1).




FIGURE 1. A terminal point, a rupture vertex and its star

Let e be a vertex of Γ . Let us denote by Γ_e the subgraph $\Gamma - \text{star}(e)$ of Γ . Since Γ is a tree, the graph Γ_e is disconnected if e is not a terminal vertex. Each connected component of Γ_e is called an *arm* of e . The number of arms of e is equal to the number of edges connected to e .

Let $e_1, \dots, e_k$ be a set of vertices of a graph Γ . The *subgraph spanned by the vertices $e_1, \dots, e_k$* is the subgraph of Γ consisting of the vertices $e_1, \dots, e_k$ and all the edges of Γ connecting any pair of vertices e_i and e_j with $1 \leq i, j \leq k$.

The complex conjugation conj of $\mathbb{C}\tilde{X}$ induced by that of $\mathbb{C}X$ defines a map on the vertices of Γ as the following:

$$\text{conj}(e_i) = \bar{e}_i = \begin{cases} e_i, & \text{if } E_i \text{ is real} \\ e_j, & \text{if } \text{conj}(E_i) = E_j. \end{cases}$$

This map is 1-1 and onto since each exceptional curve E_i is either real or it does not contain any real point at all. In fact, any normal singularity of a complex surface locally can be thought of as a branched cover of $\mathbb{C}^2$ branched along a singular curve [15]. Hence a resolution of the singularity can be constructed (as in Section 3) by taking the branched cover of a resolution of $\mathbb{C}^2$ and then by normalizing. Each exceptional curve in the resolution of $\mathbb{C}^2$ can be made real so that, in the cover, the preimage of each exceptional curve is real. This said, the claim follows (for this fact see [11] or for real resolutions see [21]).

Now, if an exceptional curve is real, the corresponding vertex will be called real; otherwise it will be called an imaginary vertex. Note that the map conj on the vertices of a resolution graph is a symmetry of the graph.

Let us denote by $\Gamma_{\mathbb{R}}$ and $\Gamma_{\mathbb{C}}$ the subgraphs spanned by the real and respectively imaginary vertices of Γ . Furthermore, let $\mathcal{C}_e$ denote the set of imaginary arms in Γ_e , that is, the arms of e on which every vertex is imaginary.

Let e be a rupture vertex of Γ . We will assign a weight n^σ to each imaginary arm σ in $\mathcal{C}_e$ and a number n'_e to vertex e . First assume that the arm $\sigma \in \mathcal{C}_e$ is a *bamboo*, i.e. contains no rupture vertex. Let there be k vertices $e_1, \dots, e_k$ on σ such that e_i is closer than e_j to e if $i < j$ ($1 \leq i, j \leq k$). Then the weight of arm σ is defined to be the continued fraction:

$$n^\sigma = n_1 - \frac{1}{n_2 - \frac{1}{\dots - \frac{1}{n_k}}}.$$

If the imaginary arms of vertex e are all bamboos, then define:

$$n'_e = n_e - \sum_{\sigma \in \mathcal{C}_e} \frac{1}{n^\sigma}.$$

Now, let an imaginary arm $\sigma \in \mathcal{C}_e$ consist of vertices $e_1, \dots, e_k$, some of which may be rupture vertices. For each rupture vertex e_i on σ compute n'_{e_i} recursively; furthermore, put $n'_i = n_i$ if e_i is a vertex on

σ that is not a rupture vertex. For a short hand notation, we put $n'_i = n'_{e_i}$. Then define

$$n^\sigma = n'_1 - \frac{1}{n'_2 - \frac{1}{\dots - \frac{1}{n'_k}}}.$$

2.3. Embedded resolution graphs. Let us consider the reduced real algebraic curve $\mathbb{C}A = \{f(x, y) = 0\}$ in $\mathbb{C}^2$ with a singularity at 0 and with no other singular point. One can resolve $\mathbb{C}A$ by a series of blow-ups of $\mathbb{C}^2$. Topologically the blown-up surface is $\mathbb{C}^2 \# t \overline{\mathbb{C}P^2}$ where t is the number of blowing-ups. Let us denote the resolution by $\rho' : \tilde{\mathbb{C}Y} \rightarrow \mathbb{C}^2$ where $\tilde{\mathbb{C}Y} = \mathbb{C}^2 \# t \overline{\mathbb{C}P^2}$. The preimage $(\rho')^{-1}(\mathbb{C}A)$ as a divisor is called the *total transform* of $\mathbb{C}A$. The closure of the preimage $(\rho')^{-1}(\mathbb{C}A - \{0\})$ as a divisor is called the *proper transform* of $\mathbb{C}A$, denoted by $\tilde{\mathbb{C}A}$. Each component of $\tilde{\mathbb{C}A}$ is smooth and can be made to transversally intersect the exceptional curves by sufficiently many blow-ups at the points of non-transversal intersection.

Let us furnish the resolution graph corresponding to the resolution ρ' with the following further data. Let the number of exceptional curves of ρ' be t . Each component of the proper transform $\tilde{\mathbb{C}A}$ which intersects an exceptional curve E_i ($1 \leq i \leq t$) is specified in the resolution graph by an arrow attached to the vertex e_i . For instance, if $f(x, y)$ is irreducible, then there exists one arrow on the graph. Furthermore, each vertex e_i is assigned a multiplicity $m_i \in \mathbb{Z}$; each arrow is assigned a multiplicity 1. Such an ‘extended’ graph is called an *embedded resolution graph* and is denoted by Γ_f . It is easy to see that Γ_f is always a tree with arrows.

We extend the definition for a rupture vertex on an embedded resolution graph as follows: A vertex of Γ_f which is adjacent to more than two vertices or two vertices and an arrow is called a *rupture vertex* of Γ_f .

The multiplicity m_i of an exceptional curve E_i is the vanishing order of the map $f \circ \rho' : \tilde{Y} \rightarrow \mathbb{C}$ on E_i . In other words, in the chart of $\tilde{Y}$ containing the intersection point of E_i and E_j , the neighbourhood of the intersection can be viewed as $\{(x, y) \in \mathbb{C}^2 | x^{m_i} y^{m_j} = 0\}$ with an appropriate choice of local coordinates (x, y) .

It is well known that the following identity is satisfied on Γ_f for every k , $1 \leq k \leq t$ (cf. e.g. [12], p. 251):

$$(3) \quad n_k m_k + \sum_{\substack{1 \leq i \leq t \\ i \neq k, n_{ik} \neq 0}} m_i = 0.$$

As a special case, it is going to be useful to understand the algorithmic construction of the embedded resolution graph for $f(x, y) = x^m + y^n$, $\gcd(m, n) = 1$. Assume that $m > n$. One can immediately observe that the blow-up sequence associated to the singularity of $f(x, y) = 0$ at 0 is directly related with the Euclidean algorithm for m and n :

$$\begin{aligned} m &= q_l n + r_l; \\ n &= q_{l-1} r_l + r_{l-1}; \\ r_l &= q_{l-2} r_{l-1} + r_{l-2}; \\ &\dots \\ r_3 &= q_1 r_2 + 1; \\ r_2 &= q_0 \cdot 1 \end{aligned}$$

where $r_l < n$, $r_i \in \mathbb{Z}^+$ and $r_{i-1} < r_i$ for all i with $1 < i < l$. The line containing q_i above corresponds to q_i consecutive blow-ups so that there are $\sum_{i=0}^l q_i$ vertices of Γ_f . Let us index the vertices of Γ_f in the following way: $e_{q_i, j}$ with $0 \leq i \leq l$ and $1 \leq j \leq q_i$ denotes the vertex created by j^{th} blow-up

within the q_i consecutive blow-ups. The unique arrow is attached to the vertex e_{q_0, q_0} . The graph Γ_f , the vertices $e_{q_i, j}$ and their multiplicities are shown in Figure 2.




FIGURE 2.

The unique rupture vertex e_{q_0, q_0} of Γ_f will be denoted by e_0 . The arm of e_0 with the terminal vertex of multiplicity n (respectively m) will be called an (n) -arm (respectively (m) -arm) of Γ_f .

3. RESOLUTION GRAPHS OF BRIESKORN DOUBLE POINTS

Let $f(x, y) = x^m + y^n$, $\mathbb{C}X$ be the double branched cover of $\mathbb{C}^2$ determined by $f(x, y) \pm z^2 = 0$, the branching locus being $\mathbb{C}A = \{f(x, y) = 0\}$. Note that the Brieskorn double surface $\mathbb{C}X$ has a normal $\mathbb{Q}$ -singularity at 0 (see e.g. [20], Chapter 1).

We are going to consider the commutative diagram:

$$\begin{array}{ccc} \mathbb{C}\tilde{X} & \xrightarrow{\rho} & \mathbb{C}X \\ \tilde{p} \downarrow & & \downarrow p \\ \mathbb{C}\hat{A} \subset \mathbb{C}\tilde{Y} & \xrightarrow{\rho'} & \mathbb{C}^2 \supset \mathbb{C}A \end{array}$$

where $p : (x, y, z) \mapsto (x, y)$ is the projection, $\rho : \mathbb{C}\tilde{X} \rightarrow \mathbb{C}X$ is a very good resolution and $\tilde{p}, \rho'$ are defined as follows.

The non-singular surface $\mathbb{C}\tilde{X}$ can be constructed in two steps. This construction, which is sketched below, can be traced, for example, in [14], Section 0; [12] Section 7.2 or [17]. At the first step, the embedded resolution graph Γ_f is constructed. Furthermore, if two exceptional curves with odd multiplicities intersect, they are made disjoint by an additional blow-up at the point of intersection. Denote the graph obtained in that way by Γ'_f and the corresponding resolution by $\rho' : \mathbb{C}\tilde{Y} = \mathbb{C}^2 \#_s \overline{\mathbb{C}P}^2 \rightarrow \mathbb{C}^2$ where s is the number of blow-ups employed so far.

Next, let $\mathbb{C}\hat{A} \subset \mathbb{C}\tilde{Y}$ be the union of the proper transform $\mathbb{C}\hat{A} = (\rho')^{-1}(\mathbb{C}A)$ and the exceptional curves in $\mathbb{C}\tilde{Y}$ with odd multiplicity. The double covering of $\mathbb{C}\tilde{Y}$ branched along $\mathbb{C}\hat{A}$ with an appropriate choice of real structure gives $\mathbb{C}\tilde{X}$. The covering map is denoted by $\tilde{p}$.

If E_i is an exceptional curve in $\mathbb{C}\tilde{Y}$ with odd multiplicity, then the preimage $\tilde{p}^{-1}(E_i)$ is connected and has self intersection $\frac{n_i}{2}$. An exceptional curve E_i of even multiplicity intersects either with two vertices of odd multiplicity or with no vertices of odd multiplicity. In case of former, $\tilde{p}^{-1}(E_i)$ is connected and has self intersection $2n_i$. In the latter case, $\tilde{p}^{-1}(E_i)$ is constituted of two exceptional curves, each with self-intersection n_i .

The graph constructed in this way is a good resolution graph and can be changed into a minimal good resolution graph by blowing-down the (-1) -curves that do not intersect more than two exceptional curves. We are going to denote by $\Gamma(m, n)$ the resolution graph corresponding to ρ .

Let the graph $\Gamma(m, n)$ have r vertices. and let E^i (respectively e^i) ($1 \leq i \leq s$) denote the exceptional curves of ρ (respectively the corresponding vertices of $\Gamma(m, n)$). The covering map $\tilde{p} : \mathbb{C}\tilde{X} \rightarrow \mathbb{C}\tilde{Y}$ induces a map $\tilde{p} : \Gamma(m, n) \rightarrow \Gamma_f$, sending e^i to e_j if E^i lies in $\tilde{p}^{-1}(E_j)$.

The following facts on Γ_f and $\Gamma(m, n)$ are direct consequences of the algorithm summarised above and at the end of previous section.

Lemma 1. Let $f(x, y) = x^m + y^n$ ($\gcd(m, n) = 1$), e_0 be the rupture vertex of Γ_f and $\tilde{p} : \Gamma(m, n) \rightarrow \Gamma_f$ be the covering map induced by $\tilde{p} : \mathbb{C}X = \{x^m + y^n \pm z^2 = 0\} \rightarrow \mathbb{C}^2 \# t\overline{\mathbb{C}P}^2$. Then:

- (i) Multiplicity of $e_0 = m_{e_0} = mn$.
- (ii) $\tilde{p}^{-1}(e_0)$ is one vertex. Let us denote it by $e^0 \in \Gamma(m, n)$.
- (iii) Let σ be an arm of e_0 . Each connected subgraph of $\Gamma(m, n)$ in the preimage $\tilde{p}^{-1}(\sigma)$ is an arm of e^0 .
- (iv) There are $\gcd(m, 2)$ arms of e^0 in the preimage of an (n) -arm of Γ_f . Analogously, there are $\gcd(n, 2)$ arms of e^0 in the preimage of an (m) -arm.
- (v) Number of arms of e^0 is 3 and each arm of e^0 is a bamboo.

The fact (iii) above allows the following definition: each arm of $\Gamma(m, n)$ in the preimage of an (n) -arm of Γ_f will be called an (n) -arm of $\Gamma(m, n)$. Analogous definition is made for an (m) -arm of $\Gamma(m, n)$.

Note that any real algebraic surface defined as a double cover of $\mathbb{C}^2$ branched along the reduced curve $\{f(x, y) = 0\}$ can be assigned two complex conjugations (see e.g. [5], Section 2.2.4). For the case of Brieskorn double surfaces when $f(x, y) = x^m + y^n$, let us denote the complex conjugations by $\text{conj}_\pm$ and the corresponding complex surfaces $\{x^m + y^n \pm z^2 = 0\}$ by $\mathbb{C}X_\pm$. First, we give a proof of the following fact pointed out in [10]:

Proposition 2. Let $\Gamma_\mathbb{R}^\pm(m, n)$ be the subgraph spanned by the vertices of $\Gamma(m, n)$ fixed under the action of $\text{conj}_\pm$. If m and n are odd, then

$$\Gamma_\mathbb{R}^\pm(m, n) = \Gamma(m, n).$$

If m is odd and n is even, then

$$\Gamma_\mathbb{R}^-(m, n) = \Gamma(m, n)$$

and $\Gamma_\mathbb{R}^+(m, n)$ is the subgraph of $\Gamma(m, n)$ spanned by the rupture vertex and the (n) -arm.

Proof. If m and n are both odd, then the resolution graph $\Gamma(m, n)$ has no symmetry but the identity. Since complex conjugations correspond to symmetries of the resolution graph, both choices of complex conjugation coincide and all exceptional curves are fixed under conjugation of either choice so that the first assertion follows.

Now let us assume that m is odd and n is even. Let $f(x, y) = x^m + y^n$ and let e_0 be the rupture vertex of Γ_f . The multiplicity m_0 of E_0 is mn and the self intersection is -1 . The embedded resolution graph Γ_f around the vertex e_0 is shown in Figure 3(a). The two vertices e_1 and e_2 adjacent to e_0 have multiplicities of different parities by Equation 3, Section 2.3. Let us assume that m_1 is odd and m_2 is even. In a properly chosen chart of $\mathbb{C}^2 \# t\overline{\mathbb{C}P}^2$ around the point of intersection $E_0 \cap E_1$, the configuration of the exceptional curves E_0, E_1, E_2 and the proper transform $\mathbb{C}\tilde{A}$ is shown in Figure 3(b). The curve $E_1 \cup \mathbb{C}\tilde{A}$ is in the branching locus and is expressed by $x^{mn}(x+1) = 0$ in this chart.



FIGURE 3.

Suppose that the double covering $\mathbb{C}X$ of $\mathbb{C}^2$ branched along $\mathbb{C}A$ is determined by $\{x^m + y^n - z^2 = 0\}$. Then above the subset $\mathcal{S} = \{(x, y) \mid -1 < x < 0\}$ of the chart in Figure 3(b) lie the imaginary points

of $\mathbb{C}X$ and above the points outside $\mathcal{S}$ on the chart lie the real points of $\mathbb{C}X$. In particular, each connected component of the preimage $\tilde{p}^{-1}(E_2)$ is real so that all the exceptional curves on its arm have real connected components on their preimages. Also $\tilde{p}^{-1}(\mathbb{C}\tilde{A})$, $\tilde{p}^{-1}(E_0)$ and $\tilde{p}^{-1}(E_1)$ are real in $\mathbb{C}X$.

Alternatively, suppose that $\mathbb{C}X = \{x^m + y^n + z^2 = 0\}$. It follows with an analogous argument to the previous paragraph that this time each connected component of the preimage $\tilde{p}^{-1}(E_2)$ contains no real point because of the sign change. $\square$

Proposition 3. *Let*

$$n' = \begin{cases} n + 2m & m \text{ odd} \\ n + m & m \text{ even.} \end{cases}$$

For $f(x, y) = x^m + y^n$ and $f'(x, y) = x^m + y^{n'}$, the resolution graph $\Gamma_{f'}$ differs from Γ_f by a pair of terminal vertices. The outermost vertex has multiplicity m and self intersection -2 while the adjacent vertex has multiplicity $2m$ (see Figure 4). Furthermore, the resolution graph $\Gamma(m, n')$ differs from $\Gamma(m, n)$ by a terminal vertex on each (n) -arm of $\Gamma(m, n')$.

Proof. Let us denote the surfaces $\{f(x, y) + z^2 = 0\}$ and $\{f'(x, y) + z^2 = 0\}$ by X and X' and the branching curves $\{f(x, y) = 0\}$ and $\{f'(x, y) = 0\}$ by A and A' respectively.




FIGURE 4. $\Gamma_{f'}$ has two extra vertices for odd m

First suppose that m is odd. Let us blow-up $\mathbb{C}^2$ once at 0 to obtain a proper transform $A'_1 \subset \mathbb{C}^2 \# \overline{\mathbb{C}P}^2$ of A' and an exceptional curve E_1 with self intersection $n_1 = -1$ and multiplicity $m_1 = m$. Then let us blow-up the $\mathbb{C}^2$ -chart of $\mathbb{C}^2 \# \overline{\mathbb{C}P}^2$ containing the singular point of A'_1 . The exceptional set contains the exceptional curve E_1 with $n_1 = -2$ transversally intersecting an exceptional curve E_2 with $n_1 = -1$, $m_2 = 2m$. Furthermore the proper transform A'_2 of A'_1 has a singularity at $A'_2 \cap E_2$ given locally by $f(x, y) = x^m + y^n = 0$, identical to the curve A .

Further blow-ups to resolve the singularity of A'_2 increase the self intersection of E_2 at least by 1 and produce a proper transform A'_t of A' and the exceptional curves $E_3, \dots, E_t$. The curves $A', E_3, \dots, E_t$ are transverse to E_2 and constitute the graph Γ_f which is a subgraph of $\Gamma_{f'}$. Therefore the embedded resolution graph $\Gamma_{f'}$ associated to f' differs from Γ_f by the two vertices e_1 and e_2 appended to one end of Γ_f .

Now consider the very good resolution $\rho' : \widetilde{X}' \rightarrow \mathbb{C}^2 \# t\overline{\mathbb{C}P}^2$ associated to $\Gamma_{f'}$. The preimages $(\rho')^{-1}(E_1)$ and $(\rho')^{-1}(E_2)$ are rational curves with self intersections -1 and $-2n_2 \leq -4$ respectively. Blowing $(\rho')^{-1}(E_1)$ down, we end up with a vertex on $\Gamma(m, n')$ with self intersection at most -3 (before further possible blow-downs of -1 curves). This vertex is appended to the (n) -arm of $\Gamma(m, n)$.

For the case of even m , the proof is analogous except that in the beginning one needs to blow-up only once to obtain a new vertex in $\Gamma(m, n')$ representing an exceptional curve of multiplicity m and self intersection $-k$. Also note that, different from the odd case, there are two (n) -arms on Γ_f and two (n') -arms on $\Gamma_{f'}$. $\square$

Corollary 4. $\Gamma(m, n + 4m/(m, 2))$ differs from $\Gamma(m, n)$ by two extra vertices at the end of each of its (n) -arms.

Proposition 5. *The self intersection of the terminal vertex of each (n) -arm of the graph $\Gamma(m, n + 4m/(m, 2))$ is -2 .*

Proof. Assume first that m is odd. Let $n' = n + 4m$ and $f'(x, y) = x^m + y^{n'}$. One has the terminal part of an (n') -arm of $\Gamma_{f'}$ as shown in Figure 5(a). To see this one needs to apply four consequent blow-ups as described in the proof of Proposition 3. Then it is straightforward to verify that the terminal part of an (n') -arm of $\Gamma(m, n')$ is as in Figure 5(b). Blowing down the (-1) -curves, the proof follows.




FIGURE 5. Resolution graphs (a) $\Gamma_{f'}$, (b) $\Gamma(m, n + 4m)$ for odd m

The case for even m is similar. □

4. THE CHARACTERISTIC SET OF A RESOLUTION GRAPH AND THE FIRST CHERN CLASS

Let us consider a very good resolution $\rho : \widetilde{\mathbb{C}U} \rightarrow \mathbb{C}U$ of a compact cone-like neighbourhood $\mathbb{C}U$ of a normal singularity and denote $\widetilde{\mathbb{C}M} = \partial\widetilde{\mathbb{C}U}$. As in [8], let c be the unique class in $H^2(\widetilde{\mathbb{C}U}, \widetilde{\mathbb{C}M}; \mathbb{Q}) \cong H^2(\widetilde{\mathbb{C}U}, \widetilde{\mathbb{C}M}) \otimes \mathbb{Q}$ such that $j^*(c) = c_1(\widetilde{\mathbb{C}U}) \in H^2(\widetilde{\mathbb{C}U}; \mathbb{Z})$ where $j : \widetilde{\mathbb{C}U} \rightarrow (\widetilde{\mathbb{C}U}, \widetilde{\mathbb{C}M})$ is the inclusion map. The canonical class $K \in H_2(\widetilde{\mathbb{C}U}; \mathbb{Q})$ is defined to be the Lefschetz dual $LD(-c)$ and can be expressed as $\sum_{1 \leq i \leq s} a_i E_i$ where $a_i \in \mathbb{Q}$ and s is the number of exceptional curves of the resolution.

If the singularity above is numerically Gorenstein, i.e. K is integral, then K is integral dual to the second Stiefel-Whitney class $w = w_2(\widetilde{\mathbb{C}U}, \widetilde{\mathbb{C}M}; \mathbb{Z})$. Therefore, we can write $w = \sum_{1 \leq i \leq s} \hat{a}_i E_i$ where $\hat{a}_i = (a_i \bmod 2)$ and E_i is identified with its dual in $H^2(\widetilde{\mathbb{C}U}, \widetilde{\mathbb{C}M}; \mathbb{Z})$. Note that w satisfies the Wu formula:

$$(4) \quad w \cdot E_i \equiv E_i \cdot E_i \pmod{2} \text{ for all } i, 1 \leq i \leq s$$

and furthermore for $1 \leq i, j \leq s$, the matrix of the intersection form $Q_{\widetilde{\mathbb{C}U}}$ on $\widetilde{\mathbb{C}U}$ has the entries

$$E_i \cdot E_j = \begin{cases} n_{ij}, & i \neq j \\ n_i, & i = j \end{cases}$$

in the basis $E_1, \dots, E_s$.

For a normal, numerically Gorenstein $\mathbb{Q}$ -singularity, let Γ be a very good resolution graph with s vertices and w be expressed as a sum as above. The *characteristic set* $\mathcal{W}$ of Γ is defined as the following set of vertices of Γ :

$$\mathcal{W} = \{e_i \in \Gamma \mid \hat{a}_i = 1\}.$$

It is algorithmic to determine the characteristic set of a very good resolution graph. Nevertheless, the relation between $c_1(\widetilde{\mathbb{C}U})$ and $\mathcal{W}$ may lead to some general properties.

Now let $p : \mathbb{C}X \rightarrow \mathbb{C}^2$ be a double covering branched along an irreducible singular curve $\mathbb{C}A = \{f(x, y) = 0\}$ that has an isolated singularity at 0. Note that the singularity of $\mathbb{C}X$ is normal and numerically Gorenstein (see e.g. [8]).

As in Section 3, consider the commutative diagram:

$$\begin{array}{ccc} \mathbb{C}\tilde{U} & \xrightarrow{\rho} & \mathbb{C}U \\ \tilde{p} \downarrow & & \downarrow p \\ \mathbb{C}\tilde{V} & \xrightarrow{\rho'} & B^4 \subset \mathbb{C}^2 \end{array}$$

where B^4 is a sufficiently small ball around 0 so that $\mathbb{C}U$ is cone-like. Let $\tilde{A}$ denote the proper transform of $\mathbb{C}A \cap B^4$ under ρ' and $\hat{A} \subset \mathbb{C}\tilde{V}$ the branching locus of the double covering $\tilde{p}$; $E^i \subset \mathbb{C}\tilde{U}$ ($1 \leq i \leq s$) denote the exceptional curves of the very good resolution ρ ; $E_i \subset \mathbb{C}\tilde{V}$ ($1 \leq i \leq t$) denote the exceptional curves of the resolution ρ' . Denote the multiplicity of E_i by m_i . We can express

$$c_1(\mathbb{C}\tilde{U}) = \sum_{i=1}^s a_i E^i, \quad c_1(\mathbb{C}\tilde{V}) = \sum_{i=1}^t b_i E_i.$$

Above and what follows below, $E^i \in H_2(\mathbb{C}\tilde{U})$ is always identified with the image $j_U^*(LD(E_i))$ of its Lefschetz dual $LD(E_i) \in H^2(\mathbb{C}\tilde{U}, \partial\mathbb{C}\tilde{U})$ under the homomorphism j_U^* induced by the inclusion $j_U : \mathbb{C}\tilde{U} \rightarrow (\mathbb{C}\tilde{U}, \partial\mathbb{C}\tilde{U})$ and E_i is identified with the image $j_V^*(LD(E_i))$ under the inclusion $j_V : \mathbb{C}\tilde{V} \rightarrow (\mathbb{C}\tilde{V}, \partial\mathbb{C}\tilde{V})$.

Proposition 6. *Assume that $E_j = \tilde{p}(E^i)$ for some i and j with $1 \leq i \leq s$ and $1 \leq j \leq t$. If the multiplicity m_j of E_j is odd then the coefficient a_i in $c_1(\mathbb{C}\tilde{U})$ is even.*

Proof. First note that $c_1(\mathbb{C}\tilde{U}) = (\tilde{p})^*(c_1(\mathbb{C}\tilde{V}) - \frac{1}{2}[\hat{A}])$ (see e.g. [2]). The class $c_1(\mathbb{C}\tilde{V}) - \frac{1}{2}[\hat{A}]$ can be expressed as a sum $\sum_{i=1}^t d_i E_i$ in $H^2(\mathbb{C}\tilde{V})$ with $d_i \in \mathbb{Z}$. Then,

$$c_1(\mathbb{C}\tilde{U}) = (\tilde{p})^*\left(\sum_{i=1}^t d_i E_i\right) = \sum_{\substack{1 \leq i \leq t \\ m_i \text{ odd}}} 2d_i (\tilde{p})^*(E_i) + \sum_{\substack{1 \leq i \leq t \\ m_i \text{ even}}} d_i (\tilde{p})^*(E_i)$$

which shows that all of the exceptional curves with odd multiplicities have even coefficients in $c_1(\mathbb{C}\tilde{U})$. $\square$

Proposition 7. *Assume that $E_j = \tilde{p}(E^i)$ for some i and j with $1 \leq i \leq s$ and $1 \leq j \leq t$. Also let $(\tilde{p})^*(\frac{1}{2}[\hat{A}]) = \sum_{k=1}^s d_k E^k$, $d_k \in \mathbb{Z}$. Then d_i is odd if and only if the multiplicity $m_j \equiv 2 \pmod{4}$.*

Proof. First let us note:

Lemma 8. $[\tilde{A}] = -\sum_{l=1}^t m_l E_l.$

Proof. The proper transform $\tilde{A}$ intersects just one exceptional curve, say E_r . To prove the claim, it is enough to check that

$$Q_{\mathbb{C}\tilde{V}}\left(-\sum_{l=1}^t m_l E_l, E_r\right) = \begin{cases} 1, & r = l \\ 0, & r \neq l. \end{cases}$$

It is then straightforward to see that this identity is in fact a reformulation of Equation 3, Section 2.3. $\square$

Since

$$[\hat{A}] = [\tilde{A}] + \sum_{m_i \text{ odd}} E_i,$$

it follows from the lemma that

$$\begin{aligned} [\widehat{A}] &= -\sum_{i=1}^t m_i E_i + \sum_{\substack{1 \leq i \leq t \\ m_i \text{ odd}}} E_i \\ &= -\sum_{\substack{1 \leq i \leq t \\ m_i \text{ even}}} m_i E_i + \sum_{\substack{1 \leq i \leq t \\ m_i \text{ odd}}} (-m_i + 1) E_i. \end{aligned}$$

Considering the behaviour of the transfer map from $H^2(\mathbb{C}\widetilde{U}; \mathbb{Z})$ to $H^2(\mathbb{C}\widetilde{V}; \mathbb{Z})$ one can deduce that

$$(\widetilde{p})^* \left(\frac{1}{2} [\widehat{A}] \right) = \sum_{\substack{1 \leq k \leq t \\ m_k \text{ even}}} \frac{m_k}{2} (\widetilde{p})^* (E_k) + \sum_{\substack{1 \leq k \leq t \\ m_k \text{ odd}}} (-m_k + 1) (\widetilde{p})^* (E_k).$$

A coefficient on the right hand side corresponding to E_j is odd if and only if $m_j \equiv 2 \pmod{4}$. $\square$

The last two propositions now prove

Proposition 9. *Let $E_j = \widetilde{p}(E^i)$ for some i and j with $1 \leq i \leq s$ and $1 \leq j \leq t$. Then a_i is odd if and only if one of the following two conditions holds: (i) $m_j \equiv 2 \pmod{4}$ and b_j even; (ii) $4|m_j$ and b_j odd.*

Proposition 10. *Corresponding to the resolution $\rho' : \mathbb{C}\widetilde{V} \rightarrow B^4$, consider the blow-up sequence:*

$$\begin{array}{ccccccccccccccc} \mathbb{C}\widetilde{V} = Y_t & \xrightarrow{\rho_t} & \dots & \xrightarrow{\rho_{k+1}} & Y_k & \xrightarrow{\rho_k} & Y_{k-1} & \xrightarrow{\rho_{k-1}} & \dots & \xrightarrow{\rho_1} & Y_0 = B^4 \subset \mathbb{C}^2 \\ \cup & & & & \cup & & \cup & & & & \cup \\ \widetilde{A} & \longrightarrow & \dots & \longrightarrow & A_k & \longrightarrow & A_{k-1} & \longrightarrow & \dots & \longrightarrow & A_0 = \mathbb{C}A \cap B^4 \end{array}$$

where Y_i ($1 \leq i \leq t$) is obtained by blowing up Y_{i-1} at the singular point of A_{i-1} , ρ_i is the corresponding map and A_i is the proper transform of A_{i-1} . Let E_k denote the exceptional sphere $Cl(\rho_k^{-1}(A_{k-1}) - A_k)$

in Y_k . Assume that $c_1(Y_k) = \sum_{i=1}^k b_i^{(k)} E_i$, $b_i^{(k)} \in \mathbb{Z}$. Then,

$$c_1(Y_{k+1}) = c_1(Y_k) + \left(-1 + \sum_{\substack{1 \leq i \leq k \\ E_i \cap E_{k+1} \neq \emptyset}} b_i^{(k)} \right) E_{k+1}.$$

Proof. It is enough to show that $c_1(Y_{k+1})$ as expressed above satisfies the adjunction formula $E_i \cdot E_i + 2 = c_1(Y_{k+1})(E_i)$ for all i , $1 \leq i \leq k+1$. Let $n_i^{(k)}$ denote the self intersection of E_i in Y_k . Then the self intersection $n_i^{(k+1)}$ of E_i in Y_{k+1} is decreased by 1 if $E_i \cap A_k \neq \emptyset$ in Y_k and is unchanged otherwise. Furthermore, $n_{k+1}^{(k+1)} = -1$ and by Equation 3, Section 2.3, $m_{k+1} = 1 + \sum_{\substack{1 \leq i \leq k \\ E_i \cap E_{k+1} \neq \emptyset}} m_i$. It is now

straightforward to check that $c_1(Y_{k+1})$ as given in the form above satisfies the adjunction formula. $\square$

Remark 11. Assume that $c_1(Y_k) = \sum_{i=1}^k d_i E_i$. It follows from the previous proposition that

$$c_1(Y_{k-1}) = \sum_{i=1}^{k-1} d_i E_i.$$

Proposition 12. *Let $\mathbb{C}A = \{f(x, y) = x^m + y^n = 0\} \cap B^4$ with $\gcd(m, n) = 1$. Assume that E_i ($1 \leq i \leq t$) are as above and that $c_1(\mathbb{C}\widetilde{V}) = \sum_{i=1}^t b_i E_i$. Then b_t is odd if and only if both m and n are odd.*

Proof. Let us adopt the notation of the remark at the end of Section 2.3 so that $t = \sum_{k=0}^l q_k$. Using Proposition 10, one can algorithmically calculate the coefficients in the expression of $c_1(\mathbb{C}\tilde{V})$. Let $b_{q_i,j}$ be the coefficient corresponding to vertex $e_{q_i,j}$. Then, $b_{q_l,1} = -1, b_{q_l,2} = -2 \dots b_{q_l,q_l} = -q_l, b_{q_{l-1},1} = -q_l - 1, b_{q_{l-1},2} = -2q_l - 2 \dots b_{q_{l-1},q_{l-1}} = -q_{l-1}q_l - q_{l-1} \dots$. Then it is straightforward to verify by induction on l that:

$$-b_t = -b_{q_0,q_0} = q_l n + q_{l-1} r_l + q_{l-2} r_{l-1} + \dots + q_1 r_2 + q_0 \cdot 1 = m + n - 1.$$

Hence the proof follows. $\square$

5. COMPUTATION OF THE INVARIANT VIA RESOLUTION GRAPHS

I give the proof of the following fact which was stated in [10].

Proposition 13. *Let $\mathbb{C}E$ be a real rational $(-m)$ -curve ($m > 0, m \in \mathbb{Q}$) in a real algebraic Q -surface $\mathbb{C}X'$. Let $\mathbb{R}E$ be smooth and $\mathbb{R}X'$ be nonorientable. Consider the projection $p : \mathbb{C}X' \rightarrow X = \mathbb{C}X'/\mathbb{C}E$. Then the point $x = p(\mathbb{C}E) \in X$ is a Q -singularity. Furthermore one can define a bilinear form tb_x on $H_1(M)$ (where $M = S_\epsilon \cap p(\mathbb{R}X')$) by pushing the framings via p as in Section 2.1. The matrix of the form tb_x in the basis formed by the connected components of M is given by:*

$$[\text{tb}_x] = \begin{cases} \begin{pmatrix} -\frac{m}{4} & -\frac{m}{4} \\ -\frac{m}{4} & -\frac{m}{4} \end{pmatrix} & , \quad \mathbb{R}E \text{ 2-sided on } \mathbb{R}X' \\ (-m) & , \quad \mathbb{R}E \text{ 1-sided on } \mathbb{R}X'. \end{cases}$$

Proof. The point x is a Q -singularity because a sufficiently small neighbourhood $\mathbb{C}U'$ of $\mathbb{C}E$ in $\mathbb{C}X'$ has a boundary which is a $\mathbb{Q}$ -homology sphere.

Let U denote $p(\mathbb{C}U')$. Let us consider the map $c' : \mathbb{C}U' \rightarrow \overline{U}$ of quotient by conjugation.

Assume first that $\mathbb{R}E$ is 2-sided on $\mathbb{R}X'$. For $i = 1, 2$ put $\gamma_i = \partial \mathbb{C}U' \cap \mathbb{R}X'$; γ_1 and γ_2 are the two connected components of a small neighbourhood of the cycle $\mathbb{R}E$ in $\mathbb{R}X'$. Let A_i be the annulus on $\mathbb{R}X'$ bounded by γ_i and $\mathbb{R}E$. Assume that the canonical orientation of the disk $\overline{E} = \mathbb{C}E/\text{conj}$ agrees with A_1 . Hence, since the surface $\mathbb{R}X'$ is nonorientable, the orientation of $\overline{E}$ also agrees with the orientation of A_2 . Now consider the topological disks $\Sigma_1 = A_1 \cup_{\mathbb{R}E} \overline{E}$ and $\Sigma_2 = A_2 \cup_{\mathbb{R}E} \overline{E}$ in $\overline{U}$. Then it follows from Equation 1, Section 2, that:

$$\overline{\text{tb}}_x(\gamma_1, \gamma_1) = \langle \Sigma_1, \Sigma_1 \rangle_{(\overline{U}, \partial \overline{U})} = \frac{1}{2} \langle \mathbb{C}E, \mathbb{C}E \rangle_{(\mathbb{C}U', \partial \mathbb{C}U')} = -\frac{m}{2};$$

$$\overline{\text{tb}}_x(\gamma_2, \gamma_2) = \langle \Sigma_2, \Sigma_2 \rangle_{(\overline{U}, \partial \overline{U})} = \frac{1}{2} \langle \mathbb{C}E, \mathbb{C}E \rangle_{(\mathbb{C}U', \partial \mathbb{C}U')} = -\frac{m}{2};$$

$$\overline{\text{tb}}_x(\gamma_1, \gamma_2) = \langle \Sigma_1, \Sigma_2 \rangle_{(\overline{U}, \partial \overline{U})} = \frac{1}{2} \langle \mathbb{C}E, \mathbb{C}E \rangle_{(\mathbb{C}U', \partial \mathbb{C}U')} = -\frac{m}{2}.$$

First part of the proof follows from Equation 2, Section 2.1.

Now assume that the cycle $\mathbb{R}E$ is 1-sided on $\mathbb{R}X'$. In this case, a sufficiently small neighbourhood of $\mathbb{R}E$ in $\mathbb{R}X'$ is a Möbius band, say A , thus has connected boundary, say γ . Observing that $\gamma = 2\mathbb{R}E$, define Σ to be the relative 2-chain $A \cup_{\mathbb{R}E} 2\overline{E}$ in $\overline{U}$. It follows that:

$$\overline{\text{tb}}_x(\gamma, \gamma) = \langle \Sigma, \Sigma \rangle_{(\overline{U}, \partial \overline{U})} = \frac{1}{2} \cdot 4 \langle \mathbb{C}E, \mathbb{C}E \rangle_{(\mathbb{C}U', \partial \mathbb{C}U')} = -2m.$$

$\square$

Corollary 14. *In the previous proposition, let $\mathbb{R}X'$ be orientable. Then*

$$\text{tb}_x = \begin{pmatrix} -\frac{m}{4} & \frac{m}{4} \\ \frac{m}{4} & -\frac{m}{4} \end{pmatrix}.$$

Proof. We should only note that in the case of orientable $\mathbb{R}X'$, $\mathbb{R}E$ is 2-sided on $\mathbb{R}X'$ and the orientation of $\overline{E}$ does not agree with the orientations of A_1 and A_2 simultaneously. $\square$

Theorem 15. *Let $x \in \mathbb{C}X$ be a normal, numerically Gorenstein, unibranch real Q -singularity; $\mathbb{C}U$ a small compact cone-like real neighbourhood of x ; $\mathbb{R}M = \mathbb{R}X \cap \partial\mathbb{C}U$; Γ be the graph of a very good resolution $\rho : \mathbb{C}\tilde{U} \rightarrow \mathbb{C}U$ at x ; $\mathbb{R}\tilde{U}$ be the preimage $\rho^{-1}(\mathbb{R}U)$; $\mathcal{W}_{\mathbb{R}}$ be the characteristic set on $\Gamma_{\mathbb{R}}$ and N be the number of vertices on $\Gamma_{\mathbb{R}}$. Then:*

$$(5) \quad \text{tb}(\mathbb{R}M) = N - 1 + \sum_{\substack{1 \leq i \leq s \\ e_i \in \mathcal{W}_{\mathbb{R}}}} n'_i.$$

Proof. First note that the normality of the singularity x implies that Γ is connected. Since x is assumed to be numerically Gorenstein, the canonical class of $\mathbb{C}\tilde{U}$ is integral so that one can define the characteristic set of the graph. Furthermore since x is a Q -singularity, the graph Γ is a tree and each exceptional curve is a sphere (see e.g. [6]).

The surface $\mathbb{R}\tilde{U}$ is smooth, connected and with boundary $\mathbb{R}M$. $\mathbb{R}\tilde{U}$ need not to be orientable. It is nonorientable if and only if n_i ($1 \leq i \leq s$) is odd for at least one real exceptional sphere (see [21], Proposition II.6.9 and Corollary II.6.10). If n_i is even for all exceptional spheres, then $\mathbb{R}\tilde{U}$ is oriented and $c_1(\mathbb{C}\tilde{U})|_{\mathbb{R}\tilde{U}}$ is even, i.e. by definition $\mathcal{W}_{\mathbb{R}}$ is empty, so that the second term on the right hand side of Equation 5 disappears and one obtains Equation 1, Section 2:

$$\text{tb}(\mathbb{R}M) = -\chi(\mathbb{R}\tilde{U}) = N - 1.$$

In the general case where $\mathbb{R}\tilde{U}$ is nonorientable, we construct a real algebraic surface $\widehat{\mathbb{C}U}$ with an oriented real part by getting rid of the terms that appear in $c_1(\mathbb{C}\tilde{U})|_{\mathbb{R}\tilde{U}}$. So, let

$$F = \{E_i | e_i \in \mathcal{W}_{\mathbb{R}}\} \bigcup \{E_i | e_i \in \mathcal{C}_j \text{ for some } j (1 \leq j \leq s), \text{ such that } e_j \in \mathcal{W}_{\mathbb{R}}\}.$$

We define:

$$\text{pr} : \mathbb{C}\tilde{U} \rightarrow \widehat{\mathbb{C}U} = \mathbb{C}\tilde{U}/F.$$

It is well-known that $c_1(\widehat{\mathbb{C}U})|_{\mathbb{R}\widehat{U}}$ is even if and only if $w_1(\mathbb{R}\widehat{U})$ is zero. Since each exceptional curve in the canonical class of $\mathbb{C}U$ is contracted in $\widehat{\mathbb{C}U}$, the real part $\mathbb{R}\widehat{U}$ is orientable. However, $\mathbb{R}\widehat{U}$ is not smooth. Let the number of singularities of $\widehat{\mathbb{C}U}$ be t which is exactly the number of vertices in $\mathcal{W}_{\mathbb{R}}$. Each singular point is real and is the image $\text{pr}(E_i)$ where E_i corresponds to the vertex $e_i \in \mathcal{W}_{\mathbb{R}}$; we denote this singular point by x_i .

Lemma 16. *Each singular point of $\widehat{\mathbb{C}U}$ is a Q -singularity.*

Proof. Let x be such a singularity. The subgraph of Γ spanned by the vertices corresponding to the exceptional spheres in the preimage $\text{pr}^{-1}(x)$ is a connected, negative definite tree, i.e. a small regular neighbourhood of $\text{pr}^{-1}(x)$ in $\mathbb{C}\tilde{U}$ has boundary a $\mathbb{Q}$ -homology sphere. $\square$

Lemma 17. *Let $x_i \in \mathbb{R}\widehat{U}$ be one of above singularities. In the basis of the homology classes of the real branches of x_i ,*

$$[\text{tb}_{x_i}] = \begin{cases} \begin{pmatrix} \frac{n'_i}{4} & \frac{n'_i}{4} \\ \frac{n'_i}{4} & \frac{n'_i}{4} \end{pmatrix} & , \text{ RE 2-sided on } \mathbb{R}X' \\ (n'_i) & , \text{ RE 1-sided on } \mathbb{R}X'. \end{cases}$$

Proof. Let $\mathbb{C}\widehat{U}_{x_i} \subset \mathbb{C}\widehat{U}$ be a sufficiently small compact cone-like neighbourhood of x_i and $\mathbb{C}M_{x_i}$ be the boundary $\partial\mathbb{C}\widehat{U}_{x_i}$. Let $\mathcal{E}$ denote the subspace of $H_2(\text{pr}^{-1}(\mathbb{C}\widehat{U}_{x_i}))$ generated by the homology classes of exceptional spheres in $\text{pr}^{-1}(x_i)$. The complex intersection form in $\mathbb{C}\widehat{U}_{x_i}$ is the restriction of the intersection form in $\text{pr}^{-1}(\mathbb{C}\widehat{U}_{x_i})$ to the orthogonal complement $\mathcal{E}^\perp$ of $\mathcal{E}$ in $H_2(\text{pr}^{-1}(\mathbb{C}\widehat{U}_{x_i}))$, since the restriction of the intersection form in $\text{pr}^{-1}(\mathbb{C}\widehat{U}_{x_i})$ to $\mathcal{E}$ is nondegenerate (see [10], Section 5.2). The result of the corresponding orthogonalisation process on the matrix of the intersection form coincides with the continued fractions of the self intersection numbers of the vertices on the contracted arms. The self intersection of $[E_i]$ restricted to the subspace $\langle \mathcal{E}^\perp, [E_i] \rangle$ is therefore equal to the number n'_i defined in Section 2.2. Since the singularity x_i is obtained by contracting E_i , the result follows from Proposition 13. $\square$

Now, since $\mathbb{R}\widehat{U}$ is oriented, by Equation 1, Section 2, we get:

$$\begin{aligned} \text{tb}(\mathbb{R}M) &= \text{tb}(\mathbb{R}\widehat{M}) = \langle \mathbb{R}\widehat{U}, \mathbb{R}\widehat{U} \rangle_{(\mathbb{C}\widehat{U}, \partial\mathbb{C}\widehat{U})} \\ &= -\chi(\mathbb{R}\widehat{U} - \{t \text{ points}\}) + \sum_{\substack{1 \leq i \leq s \\ e_i \in \mathcal{W}_{\mathbb{R}}}} \langle \Sigma_i, \Sigma_i \rangle_{(\mathbb{C}\widehat{U}_{x_i}, \partial\mathbb{C}\widehat{U}_{x_i})} \end{aligned}$$

where Σ_i is the connected piece of the surface $\mathbb{R}\widehat{U}$ bounded by the connected components of the real link $\mathbb{R}\widehat{M}_{x_i}$ of x_i (number of connected components is either one or two).

For any $e_i \in \mathcal{W}_{\mathbb{R}}$, first let the curve $\mathbb{R}E_i$ be 2-sided on the nonorientable surface $\mathbb{R}\widetilde{U}$ and γ_{i1}, γ_{i2} be the two connected components of $\mathbb{R}\widehat{M}_{x_i}$. It easy to see that

$$\begin{aligned} \langle \Sigma_i, \Sigma_i \rangle_{(\mathbb{C}\widehat{U}_{x_i}, \partial\mathbb{C}\widehat{U}_{x_i})} &= \text{tb}_{x_i}(\gamma_{i1}, \gamma_{i1}) + \text{tb}_{x_i}(\gamma_{i2}, \gamma_{i2}) + 2\text{tb}_{x_i}(\gamma_{i1}, \gamma_{i2}) \\ &= 4 \cdot \frac{n'_i}{4} = n'_i \end{aligned}$$

where the second equality follows from Lemma 17. Similarly, letting $\mathbb{R}E_i$ be 1-sided on $\mathbb{R}\widetilde{U}$, we get $\langle \Sigma_i, \Sigma_i \rangle_{(\mathbb{C}\widehat{U}_{x_i}, \partial\mathbb{C}\widehat{U}_{x_i})} = n'_i$. Hence,

$$\begin{aligned} \text{tb}(\mathbb{R}M) &= -\chi(\mathbb{R}\widehat{U} - \{t \text{ points}\}) + \sum_{\substack{1 \leq i \leq s \\ e_i \in \mathcal{W}_{\mathbb{R}}}} n'_i \\ &= -(2 - N - 1) + \sum_{\substack{1 \leq i \leq s \\ e_i \in \mathcal{W}_{\mathbb{R}}}} n'_i \\ &= N - 1 + \sum_{\substack{1 \leq i \leq s \\ e_i \in \mathcal{W}_{\mathbb{R}}}} n'_i. \end{aligned}$$

$\square$

Note that the above calculation is an application of the Integral Formula (see [9]) applied to dimension two.

Example: Consider the singularity on the surface $\mathbb{C}X_{\pm} = \{x^5 + y^8 \pm z^2 = 0\}$ with the resolution graph $\Gamma(5, 8)$ as in Figure 6. The five vertices belonging to the characteristic set $\mathcal{W}$ are marked with a cross in the figure.

The complex conjugation conj_- on the resolved surface $\mathbb{C}\tilde{X}_-$ fixes all vertices of $\Gamma(5, 8)$ (Proposition 2). Therefore $\mathcal{W}_{\mathbb{R}}$ is the set of the five vertices marked. Meanwhile, for $\mathbb{C}\tilde{X}_+$, $\mathcal{W}_{\mathbb{R}}$ contains only the rupture vertex. In both cases $\mathbb{R}\tilde{X}_{\pm}$ is nonorientable.

It follows now from Theorem 15 that

$$\text{tb}_- = 12 - 1 + 5 \cdot (-2) = 1 \text{ and } \text{tb}_+ = 2 - 1 + (-2 - 2 \cdot (-\frac{9}{11})) = \frac{7}{11}.$$

The values for $\text{tb}_{\pm}$ can be calculated by the method in [10] as well giving the same results.




FIGURE 6. Resolution graph for $x^5 + y^8 + z^2 = 0$

5.1. Computation of the linking form for singularities with more than one real branch: an example. Suppose now that the singularity $x \in \mathbb{C}X$ has ρ real branches with $\rho > 1$. Then the real part $\mathbb{R}M$ of the link of singularity is disconnected. Hence the linking form tb is given by $\rho \times \rho$ matrices. Let us take for example the complex surface $\mathbb{C}X$ defined as the double cover of $\mathbb{C}^2$ branched along $\{y(x^5 + y^4) = 0\}$. The boundary of the surface $\mathbb{R}U$ has two components α and β . With the methods of [10], one can compute that the 2×2 matrix $[\text{tb}]$ corresponding to tb in the basis $\{[\alpha], [\beta]\}$ is:

$$[\text{tb}] = \begin{pmatrix} \frac{3}{2} & -\frac{5}{2} \\ -\frac{5}{2} & \frac{3}{2} \end{pmatrix}$$

Alternatively, one can compute this matrix using the arguments in the proof of Theorem 15. The very good resolution graph Γ corresponding to the singularity at 0 is as in Figure 7 where the arrows stand for the two components of the proper transform of the branching locus.




FIGURE 7. Resolution graph for the singularity $z^2 = y(x^5 + y^4)$

There may be assigned two real structures on $\mathbb{C}\tilde{X}$. Each of the complex conjugations fixes each vertex on the resolution graph. Fix any one of the complex conjugations. Then the surface $\mathbb{R}\tilde{U}$ is a non-oriented surface with the two boundary components α and β (see Figure 8).




FIGURE 8. Real part of the surface $\{z^2 = y(x^5 + y^4)\}$ after resolution

Note that the curves $\gamma_1, \dots, \gamma_5$ separate $\mathbb{R}\tilde{U}$ into two disjoint surfaces; let us denote by Σ_1 the one which contains α in its boundary and by Σ_2 the other surface. Note that Σ_1 is orientable but Σ_2 is not.

The curves are the real parts of the -2 -spheres $E_1, \dots, E_5$. Let $\mathcal{S} = e_1, \dots, e_5$. Furthermore $\mathcal{W} = \{e_5, e_6\}$ is a characteristic set of Γ .

Similarly as in the proof of Theorem 15, let $F = \{E_1, \dots, E_5, E_6\}$ and let $\text{pr} : \mathbb{C}\tilde{U} \rightarrow \mathbb{C}\hat{U} = \mathbb{C}\tilde{U}/F$. Let us denote the $\mathbb{Q}$ -singularities $\text{pr}(E_i) \in \mathbb{C}\hat{U}$ ($i = 1, \dots, 6$) by x_i . The surface $\hat{\Sigma}_1 = \text{pr}(\Sigma_1)$ is orientable, has boundary $\text{pr}(\alpha)$ and contains five $\mathbb{Q}$ -singularities $x_1, \dots, x_5$. The surface $\hat{\Sigma}_2 = \text{pr}(\Sigma_2)$ is orientable, has boundary $\text{pr}(\beta)$ and contains six $\mathbb{Q}$ -singularities $x_1, \dots, x_5, x_6$. It follows by the proof of Theorem 15 and Proposition 13 that

$$\begin{aligned} \text{tb}(\alpha, \alpha) &= \langle \hat{\Sigma}_1, \hat{\Sigma}_1 \rangle_{\mathbb{C}\hat{U}, \partial \mathbb{C}\hat{U}} = -\chi(\hat{\Sigma}_1 - \{x_1, \dots, x_5\}) + 5 \cdot \left(-\frac{1}{2}\right) = 4 - \frac{5}{2} = \frac{3}{2}; \\ \text{tb}(\beta, \beta) &= \langle \hat{\Sigma}_2, \hat{\Sigma}_2 \rangle_{\mathbb{C}\hat{U}, \partial \mathbb{C}\hat{U}} = -\chi(\hat{\Sigma}_2 - \{x_1, \dots, x_6\}) + 5 \cdot \left(-\frac{1}{2}\right) + (-2) \\ &= 6 - \frac{5}{2} - 2 = \frac{3}{2}; \\ \text{tb}(\alpha, \beta) &= \langle \hat{\Sigma}_1, \hat{\Sigma}_2 \rangle_{\mathbb{C}\hat{U}, \partial \mathbb{C}\hat{U}} = 5 \cdot \left(-\frac{1}{2}\right) = -\frac{5}{2} = \text{tb}(\beta, \alpha). \end{aligned}$$

6. ON THURSTON-BENNEQUIN NUMBERS OF BRIESKORN DOUBLE POINTS

The Thurston-Bennequin numbers corresponding to the Brieskorn double points $x^m + y^n \pm z^2 = 0$ ($\gcd(m, n) = 1$) will be denoted by $\text{tb}_{\pm}(m, n)$.

Corollary 18. *The number $\text{tb}_{-}(m, n)$ is always an integer.*

Proof. By Proposition 2, Section 3, all vertices of $\Gamma(m, n)$ are fixed under the conj_{-} , i.e. $\Gamma_{\mathbb{R}} = \Gamma$, so that all the non-integer terms in Equation 5, Section 5, disappear. $\square$

The Thurston-Bennequin number $\text{tb}_{+}(m, n)$ is in general non-integral.

Proposition 19. *Let m be even and n odd. The rupture vertex e^0 is in $\mathcal{W}$ if and only if $4 \nmid m$.*

Proof. The vertex $e_0 = \text{conj}_{\pm}(e^0)$ in the graph Γ_f of $\mathbb{C}\tilde{Y}$ has even multiplicity mn . Therefore by Proposition 12 and Proposition 9, Section 4, the proof follows. $\square$

Corollary 20. *Assume that $4 \mid m$ and n is odd. Then $\text{tb}_{+}(m, n)$ is integer.*

Proof. From the proposition above, it follows that there is no (non-integer) contribution of imaginary arms. $\square$

The theorem below shows that if the value of $\text{tb}_{\pm}(m, n)$ is known for sufficiently small values of m and n , the number $\text{tb}_{\pm}(m', n')$ can be computed for greater integers m' and n' .

Theorem 21. *If m is odd then*

$$\text{tb}_{\pm}(m, n + 4m) = \text{tb}_{\pm}(m, n).$$

If $4|m$ then

$$\text{tb}_{\pm}(m, n + 2m) = \text{tb}_{\pm}(m, n).$$

If $m \equiv 2 \pmod{4}$ then

$$\begin{aligned} \text{tb}_{-}(m, n + 2m) - \text{tb}_{-}(m, n) &= 4, \\ \text{tb}_{+}(m, n + 2m) - \text{tb}_{+}(m, n) &= \frac{4}{n(n + 2m)}. \end{aligned}$$

Proof. Assume first that m is odd. By Corollary 4, Section 3, there are two vertices appended to each (n) -arm of $\Gamma(m, n)$ to construct $\Gamma(m, n + 4m/(m, 2))$; let us call them e_1 and e_2 with e_1 being the outer-most vertex. Take an (n) -arm of $\Gamma(m, n)$; let us call the two vertices to be appended as e_1 and e_2 with e_1 being the outer-most vertex. By Proposition 5, Section 3, e_1 has self intersection -2 .

To prove the theorem, it is enough to show that $e_1 \in \mathcal{W}$ and $e_2 \notin \mathcal{W}$ since in that case it follows from Theorem 15, Section 5, that

$$\text{tb}_{-}(m, n + 4m) = \text{tb}_{-}(m, n) - (-2) + (-2) = \text{tb}_{-}(m, n).$$

Recall that by Proposition 5, the terminal part added to Γ_f to construct the graph $\Gamma_{f'}$ corresponding to $\tilde{Y}'$ is as in Figure 5(a). The exceptional curves E'_1 and E'_2 have odd multiplicities m and $3m$ respectively. Therefore by Proposition 6, Section 4, the vertices of $\Gamma(m, n + 4m)$ corresponding to the exceptional curves $\tilde{p}^{-1}(E'_1)$ and $\tilde{p}^{-1}(E'_2)$ are not in $\mathcal{W}$. It follows that the vertex corresponding to the exceptional curve $\tilde{p}^{-1}(E_1)$ must appear in $\mathcal{W}$ for Equation 4, Section 4, to be satisfied in $\mathbb{C}\tilde{U}'$. Again in order to satisfy Equation 4, the vertex corresponding to the exceptional curve E_2 is not included in $\mathcal{W}$.

Assume now that m is divisible by 4 and n is odd. By Proposition 2, Corollary 4, Section 3, and Proposition 19, it follows that $\text{tb}_{+}(m, n + 2m) = \text{tb}_{+}(m, n)$. For the relation on the invariant tb_{-} , we first note that, as in the proof of Proposition 3, Section 3, the terminal part of the resolution graph $\Gamma_{f'}$ is as shown in Figure 9. The multiplicity of the sphere E_1 is m hence by Lemma 7, the preimage $(\tilde{p})^*(E_1)$ does not contribute mod 2 to $(\tilde{p})^*(\frac{1}{2}[\widehat{A'}])$.




FIGURE 9.

The coefficient b_1 of the cycle E_1 in $c_1(\mathbb{C}\tilde{V}')$ is -1 because E_1 is produced after the first blow-up of $\mathbb{C}^2$ (cf. Proposition 10, Section 4). Therefore by Proposition 7, Section 4, the vertex of $\Gamma(m, n + 4m)$ corresponding to $\tilde{p}^{-1}(E_1)$ is in $\mathcal{W}$. As a consequence, $\tilde{p}^{-1}(E_2)$ is not in $\mathcal{W}$.

(One can also show that $\tilde{p}^{-1}(E_2)$ is not in $\mathcal{W}$ by the following reasoning: the multiplicity of E_2 is $2m$. Since E_2 is produced after the second blow-up of $\mathbb{C}^2$, the coefficient b_2 is -2 so that $a_2 = 0 \pmod{2}$. Therefore the vertex of $\Gamma(m, n + 4m)$ corresponding to $\tilde{p}^{-1}(E_2)$ is not in $\mathcal{W}$.)

Now assume that $m \equiv 2 \pmod{4}$ and n is odd. The terminal part of the resolution graph $\Gamma_{f'}$ is again as in Figure 9. Note that $b_1 = -1$ and $b_2 = -2$ so that by Proposition 9, $e_1, e_2 \notin \mathcal{W}$. It follows from Theorem 15 that

$$\text{tb}_{-}(m, n + 2m) - \text{tb}_{-}(m, n) = -\chi(\mathbb{R}X') + \chi(\mathbb{R}X) = 2 \cdot 2 = 4.$$

For the last claim in the theorem, we observe by Proposition 19 that $e^0 \in \mathcal{W}_{\mathbb{R}}$. It follows from Corollary 4 that $\Gamma(m, n+2m)$ and $\Gamma(m, n)$ differ in the pair of $(n+2m)$ -arms of the former and pair of (n) -arms of the latter. Let σ' be an $(n+2m)$ -arm of $\Gamma(m, n+2m)$ and σ be an (n) -arm of $\Gamma(m, n)$. By Theorem 15

$$tb_+(m, n+2m) - tb_+(m, n) = 2\left(\frac{1}{n^{\sigma'}} - \frac{1}{n^{\sigma}}\right).$$

By [18] Theorem 3.6.1, it follows that

$$\frac{1}{n^{\sigma'}} - \frac{1}{n^{\sigma}} = -\frac{1}{m(n+2m)} + \frac{1}{mn} = \frac{2}{n(n+2m)}$$

and the proof follows. $\square$

Theorem 22. *Let m be odd and $k, t \in \mathbb{Z}^+$, $4km > t$. Then*

$$tb_{\pm}(m, 4km - t) + tb_{\pm}(m, t) = -2.$$

Proof. Consider the weighted homogeneous surface $\mathbb{C}P(4k, 1, 1)$ with a unique singularity at $[1 : 0 : 0]$ (cf. [6] Appendix B for a brief introduction to weighted projective spaces) and the curve $\mathbb{C}A = \{x^m + u^{4km-t}v^t = 0\}$ in $\mathbb{C}P(4k, 1, 1)$. $\mathbb{C}A$ has a pair of singularities at $[0 : 0 : 1]$ and $[0 : 1 : 0]$. Let $\mathbb{C}\tilde{P}(4k, 1, 1)$ obtained by blowing-up $\mathbb{C}P(4k, 1, 1)$ at $[1 : 0 : 0]$. Denote the resolution by ρ .

The plan of the proof is as the following. First we construct a branched double covering $p : \mathbb{C}X \rightarrow \mathbb{C}P(4k, 1, 1)$ whose branching locus contains the curve $\mathbb{C}A$. The surface $\mathbb{C}X$ has the real Brieskorn double points corresponding to $(m, t, 2)$ and $(m, 4km - t, 2)$. Then we show that $\mathbb{R}X$ is orientable and $\overline{X} = \mathbb{C}X/\text{conj}$ is a $\mathbb{Q}$ -homology sphere so that $\langle \mathbb{R}X, \mathbb{R}X \rangle_{\overline{X}} = 0$. Using this identity, we obtain a relation between $tb_{\pm}(m, 4km - t)$ and $tb_{\pm}(m, t)$.


The smooth surface $\mathbb{C}\tilde{P}(4k, 1, 1)$ is biholomorphic to the Hirzebruch surface $\mathbb{F}_{4k}$ and the section $S_{\infty} \subset \mathbb{C}\tilde{P}(4k, 1, 1)$ has self intersection $-4k$ (see e.g. [2]). Let S_0 be the zero section and F be a fiber of ruling in $\mathbb{C}\tilde{P}(4k, 1, 1)$. One can show that $[\rho^{-1}(\mathbb{C}A)] = m[S_0]$ and $([\rho^{-1}(\mathbb{C}A)] + [S_{\infty}]) = (m+1)[S_0] - 4k[F]$. Hence, $[\rho^{-1}(\mathbb{C}A)]$ is odd and $([\rho^{-1}(\mathbb{C}A)] + [S_{\infty}])$ is even. Therefore there exist a double covering $\tilde{p} : \mathbb{C}\tilde{X} \rightarrow \mathbb{C}\tilde{P}$ branched along $\rho^{-1}(\mathbb{C}A) \cup S_{\infty}$. The covering $\tilde{p}$ is associated either with the complex conjugation conj_- or with conj_+ . Let us assume that $\tilde{p}$ is associated with conj_- first.

Consider the commutative diagram:

$$\begin{array}{ccccc} \mathbb{C}\tilde{X} & \xrightarrow{\tilde{p}} & \mathbb{C}X & \xrightarrow{c'} & \mathbb{C}X/\text{conj} = \overline{X} \\ \tilde{p} \downarrow & & \downarrow p & & \downarrow \overline{p} \\ \mathbb{F}_{4k} = \mathbb{C}\tilde{P}(4k, 1, 1) & \xrightarrow{\rho} & \mathbb{C}P(4k, 1, 1) & \xrightarrow{c} & \mathbb{C}P(4k, 1, 1)/\text{conj} = \overline{P} \end{array}$$

where $p : \mathbb{C}X \rightarrow \mathbb{C}P(4k, 1, 1)$ is the double covering with branching locus $\mathbb{C}A \cup \rho(S_{\infty}) = \mathbb{C}A \cup \{[1 : 0 : 0]\} \subset \mathbb{C}P(4k, 1, 1)$ and c, c' are maps of quotient by conjugation. Note that $\mathbb{C}X = \mathbb{C}\tilde{X}/E$ where $E = \tilde{p}^{-1}(S_{\infty})$ with $E^2 = -2$ and $\mathbb{C}X$ has isolated singularities at $p^{-1}([1 : 0 : 0])$, $p^{-1}([0 : 1 : 0])$ and $p^{-1}([0 : 0 : 1])$, the last two being Brieskorn double points corresponding to $(m, t, 2)$ and $(m, 4km - t, 2)$ respectively.

It is easy to check that $c_1(\mathbb{C}\tilde{X})$ is even. Therefore the surface $\mathbb{R}\tilde{X}$ is orientable. In particular, $\mathbb{R}\tilde{X}$ is a torus and $\mathbb{R}X$ is a pinched torus. To see this, let Δ be the subset of $\mathbb{R}\tilde{P}(4k, 1, 1)$ such that $\Delta = \tilde{p}(\mathbb{R}\tilde{X})$. Then Δ is the closure of one of the connected components of $\mathbb{R}\tilde{P}(4k, 1, 1) - (\rho^{-1}(\mathbb{R}A) \cup S_{\infty})$. Therefore Δ is an annulus (see Figure 10). Note that the choice of Δ is determined by the choice of the complex conjugation $\text{conj}_{\pm}$.


 FIGURE 10. $\mathbb{R}\tilde{A} \cup \mathbb{R}S_\infty$ bounds an annulus on $\mathbb{R}\tilde{P}(4, 1, 1)$

$\mathbb{R}X$ has three singular points: $P_1 = p^{-1}([0 : 1 : 0])$, $P_2 = p^{-1}([0 : 0 : 1])$ and $P_3 = p^{-1}([1 : 0 : 0])$. The point P_3 is obtained by contracting E in $\mathbb{C}\tilde{X}$. It follows from Corollary 14, Section 5, that

$$(6) \quad [\text{tb}_{P_3}] = \begin{bmatrix} -\frac{k}{2} & \frac{k}{2} \\ \frac{k}{2} & -\frac{k}{2} \end{bmatrix}.$$

Finally we observe that $\overline{P}$ and $\overline{X}$ are $\mathbb{Q}$ -homology 4-spheres. For the former, we note the well-known fact that $\mathbb{C}P(k, 1, 1)$ is a $\mathbb{Q}$ -homology manifold with cohomology $H^*(\mathbb{C}P(k, 1, 1); \mathbb{Q}) = H^*(\mathbb{C}P^2; \mathbb{Z}) \otimes \mathbb{Q}$. Then it follows from [4], Section III.5, that $\overline{P}(k, 1, 1)$ is a $\mathbb{Q}$ -homology sphere.

In order to prove that $\overline{X}$ is a $\mathbb{Q}$ -homology sphere, it is enough to show that $\overline{p}: \overline{X} \rightarrow \overline{P}$ is branched along a 2-sphere. In fact, the branching locus is the *Arnold surface* $\mathcal{A} = \overline{A} \cup_{\partial} c(\mathbb{R}P(4k, 1, 1) - \text{int}(p(\mathbb{R}X))) \subset \overline{P}(k, 1, 1)$ where $\overline{A} = c(\mathbb{C}A)$. Since $\mathbb{C}A$ is a sphere, it follows immediately that $\mathcal{A}$ is a sphere.

Therefore there is no rational 2-homology in $\overline{X}$. In particular, the pinched torus $c'(\mathbb{R}X)$ has self intersection 0 in $\overline{X}$. It follows from this fact and Equation 6 that:

$$\begin{aligned} 0 &= \langle c'(\mathbb{R}X), c'(\mathbb{R}X) \rangle_{\overline{X}} \\ &= \text{tb}_{P_1}(\mathbb{R}X, \mathbb{R}X) + \text{tb}_{P_2}(\mathbb{R}X, \mathbb{R}X) + \text{tb}_{P_3}(\mathbb{R}X, \mathbb{R}X) - 2\chi(\mathbb{R}X - 3 \text{ points}) \\ &= \overline{\text{tb}}_{P_1} + \overline{\text{tb}}_{P_2} + \left(-\frac{k}{2} + \frac{k}{2} - \frac{k}{2} + \frac{k}{2}\right) - 2 \cdot (-2) \\ &= \overline{\text{tb}}_{P_1} + \overline{\text{tb}}_{P_2} + 4. \end{aligned}$$

The proof follows from Equation 2, Section 2.3. For an extended discussion of this proof, see also [19]. $\square$

Remark 23. *Theorem 22 is not true for the two singularities $x^m + y^{2km-t} + z^2 = 0$ and $x^m + y^t + z^2 = 0$ for arbitrary k with $k, t \in \mathbb{Z}^+$. In fact, the proof fails for odd k since in that case $\tilde{p}^{-1}(S_\infty)$ would have self intersection $\frac{-2k}{2} = -k$ in $\mathbb{C}\tilde{X}$, that is, $\mathbb{R}\tilde{X}$ and $\mathbb{R}X$ would be nonorientable so that the Equation 1 would not be valid.*

Theorem 24. *Let m be even and $k, t \in \mathbb{Z}^+$, $2km > t$. Then*

$$\text{tb}_-(m, 2km - t) + \text{tb}_-(m, t) = \begin{cases} -4, & 4|m \\ -4 + 4k, & m \equiv 2 \pmod{4}. \end{cases}$$

Proof. The idea and constructions are much similar to the previous proof. This time let us consider $\mathbb{C}A = \{x^m + u^{2km-t}v^t = 0\}$ in the weighted homogeneous surface $\mathbb{C}P(2k, 1, 1)$. The surface $\mathbb{R}P(2k, 1, 1)$ and $\mathbb{R}A$ are now as in Figure 11. Let us denote by Δ the disc bounded by $\mathbb{R}A$ in $\mathbb{R}P(2k, 1, 1)$. The curve

$\mathbb{C}A$ is even in $H_2(\mathbb{C}P(2k, 1, 1); \mathbb{Z})$ and there exists a double covering of $\mathbb{C}P(2k, 1, 1)$ with branching locus $\mathbb{C}A$.




FIGURE 11. $\mathbb{R}A$ bounds a disc on $\mathbb{R}P(2k, 1, 1)$

Let us first consider the case when the double covering is equipped with the complex conjugation conj_- ; denote the covering surface by $\mathbb{C}X_-$. Then $p(\mathbb{R}X_-) = \mathbb{R}P(2k, 1, 1) - \text{int}(\Delta)$. Let us denote by P_1 and P_2 the two singular points of $\mathbb{R}X_-$ over the two singular points of $\mathbb{C}A$. Since

$$\begin{aligned} c_1(\mathbb{C}X_-) &= p^*(c_1(\mathbb{C}P(2, 1, 1)) + \tfrac{1}{2}[\mathbb{C}A]) \\ &= (2 + \tfrac{m}{2})p^*(S_\infty) + (2(k+1) + mk)p^*(F), \end{aligned}$$

$\mathbb{R}\tilde{X}_-$ is orientable if and only if $4|m$.

The complex surface $\bar{P} = \mathbb{C}P(2k, 1, 1)/\text{conj}$ is a $\mathbb{Q}$ -homology sphere. Since the *Arnold surface* $\mathcal{A} = \bar{A} \cup_{\partial} c(\Delta)$ is a 2-sphere, it follows that $\bar{X}_- = \mathbb{C}X_-/\text{conj}$ is a $\mathbb{Q}$ -homology 4-sphere.

If $4|m$, the surface $\mathbb{R}X_-$ is an oriented surface of genus 2 with two of its nontrivial cycles pinched (see Figure 12); the two points P_3 and P_4 obtained in this way are $\mathbb{Q}$ -singularities with

$$[\text{tb}_{P_3}] = [\text{tb}_{P_4}] = \begin{bmatrix} -\frac{2k}{4} & \frac{2k}{4} \\ \frac{2k}{4} & -\frac{2k}{4} \end{bmatrix}$$

observed by Corollary 14, Section 5.




FIGURE 12.

$$\begin{aligned} \text{Hence,} \quad 0 &= \langle c'(\mathbb{R}X_-), c'(\mathbb{R}X_-) \rangle_{\bar{X}_-} \\ &= \sum_{i=1}^4 \overline{\text{tb}}_{P_i}(\mathbb{R}X_-, \mathbb{R}X_-) - 2\chi(\mathbb{R}X_- - 4 \text{ points}) \\ &= \overline{\text{tb}}_{P_1} + \overline{\text{tb}}_{P_2} + 0 + 0 + -2 \cdot (-2 - 2) \\ &= \overline{\text{tb}}_{P_1} + \overline{\text{tb}}_{P_2} + 8. \end{aligned}$$

The proof for the first claim now follows from Equation 2, Section 2.1.

If $m \equiv 2 \pmod{4}$, then $\mathbb{R}\tilde{X}_-$ is not orientable. But as in the proof of Theorem 15, we observe that $\mathbb{R}X_- = \mathbb{R}\tilde{X}_-/\tilde{p}^{-1}(S_\infty)$ is orientable. The two Q-singularities P_3 and P_4 obtained in this way have

$$[\text{tb}_{P_3}] = [\text{tb}_{P_4}] = \begin{bmatrix} -\frac{2k}{4} & -\frac{2k}{4} \\ -\frac{2k}{4} & -\frac{2k}{4} \end{bmatrix}$$

so that

$$\begin{aligned} 0 &= \sum_{i=1}^4 \overline{\text{tb}}_{P_i}(\mathbb{R}X_-, \mathbb{R}X_-) - 2\chi(\mathbb{R}X_- - 4 \text{ points}) \\ &= \overline{\text{tb}}_{P_1} + \overline{\text{tb}}_{P_2} + 2\left(-\frac{k}{2} - \frac{k}{2} - \frac{k}{2} - \frac{k}{2}\right) + -2 \cdot (-2 - 2) \\ &= \text{tb}_{P_1} + \text{tb}_{P_2} - 8k + 8 \end{aligned}$$

completing the proof for the second claim. $\square$

REFERENCES

- [1] A'Campo, N.: Le groupe de monodromie du d ploiement des singularit s isol es de courbes planes. I. (French) Math. Ann. **213**, 1–32 (1975)
- [2] Beauville A.: Surface Alg briques Complexes, Ast risque 54, Soci t  Math matique de France 1978
- [3] Bennequin D.: Entrelacements et  quations de Pfaff. Ast risque **107-108**, 87–161 (1983)
- [4] Bredon, G.E: Introduction to compact transformation groups, Pure and Applied Mathematics, Vol. 46, New York-London: Academic Press 1972
- [5] Degtyarev A.I., Kharlamov V.M.: Topological properties of real algebraic varieties: du cot  de chez Rokhlin. Russ. Math. Surv. **55** (4), 735-814 (2000)
- [6] Dimca A.: Singularities and Topology of Hypersurfaces, Springer-Verlag 1992
- [7] Durfee A.: Fifteen characterization of rational double points and simple critical points. Enseign. Math. (2) **25**, no. 1-2, 131–163 (1979)
- [8] Durfee A.: The signature of smoothings of complex surface singularities. Math. Ann. **232**, 85–98 (1978)
- [9] Finashin, S. M.: An integral formula for the complex intersection number of real cycles in a real algebraic variety with topologically rational singularities. (English. English, Russian summary) Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **279**, Geom. i Topol. **6**, 241–245, 250–251 (2001)
- [10] Finashin S. M.: Complex intersection of real cycles in real algebraic varieties and generalized Arnold-Viro inequalities. preprint, math.AG/9902022
- [11] Finashin S. M.: Rokhlin's question and smooth quotients by complex conjugation of singular real algebraic surfaces. Topology, ergodic theory, real algebraic geometry, 109–119, Amer. Math. Soc. Transl. Ser. 2, 202, Amer. Math. Soc., Providence, RI 2001.
- [12] Gompf R.E., Stipsicz A.I.: 4-Manifolds and Kirby Calculus, Graduate Studies in Mathematics 20, AMS 1999
- [13] Gusein-Zade, S.M. Intersection matrices for certain singularities of functions of two variables. (Russian, English) Funct. Anal. Appl. **8**, 10-13 (1974); translation from Funkts. Anal. Prilozh. **8**, No.1, 11-15 (1974)
- [14] Harer J., Kas A., Kirby R.: Handlebody decomposition of complex surface, Memoirs AMS, No. 350, Providence, RI, USA, vol. 62 1986
- [15] Laufer H.B.: Normal Two-Dimensional Singularities, Annals of Math. Studies 71, Princeton University Press 1971
- [16] Milnor J.: Singular Points of Complex Hypersurfaces, Annals of Math. Studies 61, Princeton University Press 1968
- [17] N methi A.: Five lectures on normal surface singularities. With the assistance of  gnes Szil rd and S ndor Kov cs. Bolyai Soc. Math. Stud., **8**, Low Dimensional Topology(Eger, 1996/Budapest, 1998), 269–351, J nos Bolyai Math. Soc., Budapest (1999)
- [18] Orlik P., Wagreich P.: Isolated singularities of algebraic surfaces with C^* action. Ann. of Math. **2 93**, 205–228 (1971)
- [19]  zt rk F.: On Thurston-Bennequin numbers of Brieskorn double points. Proceedings of Istanbul Singularity Workshop, June 2001, to appear
- [20] Saveliev, N.: Invariants for homology 3-spheres, Encyclopaedia of Mathematical Sciences, 140, Low-Dimensional Topology, I. Berlin: Springer-Verlag 2002
- [21] Silhol R.: Real Algebraic Surfaces, Lecture Notes in Math. 1392, Berlin: Springer-Verlag 1989
- [22] Varcenko A. N.: Contact structures and isolated singularities. (Russian) Vestnik Moskov. Univ. Ser. I Mat. Mekh. **101**, no. 2, 18–21 (1980)

DEPARTMENT OF MATHEMATICS, BO AZI I UNIVERSITY, İSTANBUL, TURKEY
ferit.ozturk@boun.edu.tr